

Datasheet for ABIN3301417

Human HIST1H3J cDNA Clone in Mammalian Expression Vector

Overview

Quantity:	10 µg
Gene:	HIST1H3J
Species:	Human
Insert:	cDNA
Vector:	Mammalian Expression Vector
Application:	Protein Expression (PEXP)

Product Details

Purpose:	Untagged full-length cDNA clone from Human HIST1H3J is ideal for over-expression of native protein for functional studies.
Brand:	TrueClones®
Vector Backbone:	pCMV6-Entry
Promoter:	Enhanced CMV Promoter
Selectable Marker:	Neomycin
Bacterial Resistance:	Kanamycin
Expression Type:	Transient
Specificity:	With the native stop codon at the end of the ORF the C-terminal Myc-DDK tag in the vector won't be expressed.
Characteristics:	<ul style="list-style-type: none"> • These cDNA clones are isolated from full-length cDNA libraries and usually contain the coding sequence as well as the untranslated regions (UTRs) of the mRNA transcript appropriate to the library from which they were isolated. • These cDNA clones are ideal for over-expression of native proteins for functional studies. Provided as 10 µg transfection-ready plasmids. • Every lot of primer is tested to provide clean sequencing of cDNA clones.

Order at www.genomics-online.com

USA & Canada: +1 877 302 8632 | support@antibodies-online.com

Product Details

Purification:	The DNAs were purified using PowerPrep HP Plasmid isolation kits for transfection ready plasmids.
Sequencing Primer:	VP1.5 (forward) 5'GGACTTTCCTCAAATGTTCG 3', XL39 (reverse) 5'ATTAGGACAAGGCTGGTGGG 3'
Components:	<ul style="list-style-type: none">• The cDNA clone is shipped in a 2-D bar-coded Matrix tube as dried plasmid DNA.• The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.

Target Details

Gene:	HIST1H3J
Alternative Name:	HIST1H3J (HIST1H3J Products)
Background:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015].
NCBI Accession:	NM_003535 , NP_003526

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Storage:	RT, -20 °C
Storage Comment:	The lyophilized plasmid is stable for up to one year when stored at ambient temperature. Following dissolution in 100 µL dH ₂ O, store at -20 °C. Lyophilized primers are stable for up to one year when stored at ambient temperature. Following dissolution in 10 µL dH ₂ O, store at -20 °C.

Order at www.genomics-online.com

USA & Canada: +1 877 302 8632 | support@antibodies-online.com

Page 2/3 | Product datasheet for ABIN3301417 | 09/13/2023 | Copyright antibodies-online. All rights reserved.

Handling

Expiry Date: 12 months

Publications

Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (1991)