

Datasheet for ABIN3302975

Human NFE4 cDNA Clone in Mammalian Expression Vector (Myc-DYKDDDDK Tag)

Overview

Quantity:	10 µg
Gene:	NFE4
Species:	Human
Insert:	cDNA
Vector:	Mammalian Expression Vector
Application:	Protein Expression (PEXP)

Product Details

Purpose:	Untagged full-length cDNA clone from Human NFE4 is ideal for over-expression of native protein for functional studies.
Brand:	TrueClones®
Vector Backbone:	pCMV6-Entry
Promoter:	Enhanced CMV Promoter
Selectable Marker:	Neomycin
Bacterial Resistance:	Kanamycin
Expression Type:	Transient
Specificity:	With the native stop codon at the end of the ORF the C-terminal Myc-DDK tag in the vector won't be expressed.
Characteristics:	<ul style="list-style-type: none"> • These cDNA clones are isolated from full-length cDNA libraries and usually contain the coding sequence as well as the untranslated regions (UTRs) of the mRNA transcript appropriate to the library from which they were isolated. • These cDNA clones are ideal for over-expression of native proteins for functional studies. Provided as 10 µg transfection-ready plasmids.

Order at www.genomics-online.com

USA & Canada: +1 877 302 8632 | support@antibodies-online.com

Product Details

- Every lot of primer is tested to provide clean sequencing of cDNA clones.

Purification:	The DNAs were purified using PowerPrep HP Plasmid isolation kits for transfection ready plasmids.
Sequencing Primer:	VP1.5 (forward) 5'GGACTTTCCAAAATGTCTG 3', XL39 (reverse) 5'ATTAGGACAAGGCTGGTGGG 3'
Components:	<ul style="list-style-type: none">• The cDNA clone is shipped in a 2-D bar-coded Matrix tube as dried plasmid DNA.• The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.

Target Details

Gene:	NFE4
Alternative Name:	NFE4 (NFE4 Products)
Background:	The erythroid-specific protein encoded by this gene, and the ubiquitous transcription factor CP2, form the stage selector protein (SSP) complex, which is involved in preferential expression of the gamma-globin genes in fetal erythroid cells. Alternate use of an in-frame upstream non-AUG (CUG) translation initiation codon, and a downstream AUG codon, results in two isoforms. While the long isoform (22 kDa) acts as an activator, the short isoform (14 kDa) has been shown to repress gamma-globin gene expression. This gene is located in an intron of the FBXL13 gene on the opposite strand. [provided by RefSeq, Jul 2008].
NCBI Accession:	NM_001085386 , NP_001078855

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Storage:	RT, -20 °C
Storage Comment:	The lyophilized plasmid is stable for up to one year when stored at ambient temperature. Following dissolution in 100 µL dH ₂ O, store at -20 °C. Lyophilized primers are stable for up to one year when stored at ambient temperature. Following dissolution in 10 µL dH ₂ O, store at -20 °C.

Order at www.genomics-online.com

USA & Canada: +1 877 302 8632 | support@antibodies-online.com

Handling

Expiry Date: 12 months

Publications

Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (1991)