-online.com genomics





Human NLRP8 cDNA Clone in Mammalian Expression Vector

Overview		
Quantity:	10 μg	
Gene:	NALP8 (NLRP8)	
Species:	Human	
Insert:	cDNA	
Vector:	Mammalian Expression Vector	
Application:	Protein Expression (PExp)	
Product Details		
Purpose:	Untagged full-length cDNA clone from Human NLRP8 is ideal for over-expression of native protein for functional studies.	
Brand:	TrueClones®	
Vector Backbone:	pCMV6-Entry	
Promoter:	Enhanced CMV Promoter	
Selectable Marker:	Neomycin	
Bacterial Resistance:	Kanamycin	
Expression Type:	Transient	
Specificity:	With the native stop codon at the end of the ORF the C-terminal Myc-DDK tag in the vector won't be expressed.	
Characteristics:	 These cDNA clones are isolated from full-length cDNA libraries and usually contain the coding sequence as well as the untranslated regions (UTRs) of the mRNA transcript appropriate to the library from which they were isolated. These cDNA clones are ideal for over-expression of native proteins for functional studies. Provided as 10 µg transfection-ready plasmids. Every lot of primer is tested to provide clean sequencing of cDNA clones. 	

Product Details Purification: The DNAs were purified using PowerPrep HP Plasmid isolation kits for transfection ready plasmids. Sequencing Primer: VP1.5 (forward) 5'GGACTTTCCAAAATGTCG 3', XL39 (reverse) 5'ATTAGGACAAGGCTGGTGGG • The cDNA clone is shipped in a 2-D bar-coded Matrix tube as dried plasmid DNA. Components: · The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials. **Target Details** Gene: NALP8 (NLRP8) Alternative Name: NLRP8 (NLRP8 Products) Background: NALPs are cytoplasmic proteins that form a subfamily within the larger CATERPILLER protein family. Most short NALPs, such as NALP8, have an N-terminal pyrin (MEFV, MIM 608107) domain (PYD), followed by a NACHT domain, a NACHT-associated domain (NAD), and a Cterminal leucine-rich repeat (LRR) region. The long NALP, NALP1 (MIM 606636), also has a Cterminal extension containing a function to find domain (FIIND) and a caspase recruitment domain (CARD). NALPs are implicated in the activation of proinflammatory caspases (e.g., CASP1, MIM 147678) via their involvement in multiprotein complexes called inflammasomes (Tschopp et al., 2003 [PubMed 12563287]).[supplied by OMIM, Mar 2008]. NCBI Accession: NM_176811, NP_789781 **Application Details** Restrictions: For Research Use only Handling Format: Lyophilized RT,-20 °C Storage: Storage Comment: The lyophilized plasmid is stable for up to one year when stored at ambient temperature. Following dissolution in 100 µL dH20, store at -20 °C. Lyophilized primers are stable for up to one year when stored at ambient temperature. Following dissolution in 10 µL dH2O, store at -20 °C.

12 months

Expiry Date:

Publications

Product	cited	in:

Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (1991)