

Datasheet for ABIN3322596

## Human KIR3DP1 cDNA Clone in Mammalian Expression Vector

### Overview

Quantity:	10 µg
Gene:	KIR3DP1
Species:	Human
Insert:	cDNA
Vector:	Mammalian Expression Vector
Application:	Protein Expression (PEXP)

### Product Details

Purpose:	Untagged full-length cDNA clone from Human KIR3DP1 is ideal for over-expression of native protein for functional studies.
Brand:	TrueClones®
Vector Backbone:	pCMV6-Entry
Promoter:	Enhanced CMV Promoter
Selectable Marker:	Neomycin
Bacterial Resistance:	Kanamycin
Expression Type:	Transient
Specificity:	With the native stop codon at the end of the ORF the C-terminal Myc-DDK tag in the vector won't be expressed.
Characteristics:	<ul style="list-style-type: none"> <li>• These cDNA clones are isolated from full-length cDNA libraries and usually contain the coding sequence as well as the untranslated regions (UTRs) of the mRNA transcript appropriate to the library from which they were isolated.</li> <li>• These cDNA clones are ideal for over-expression of native proteins for functional studies. Provided as 10 µg transfection-ready plasmids.</li> <li>• Every lot of primer is tested to provide clean sequencing of cDNA clones.</li> </ul>

Order at [www.genomics-online.com](http://www.genomics-online.com)

USA & Canada: +1 877 302 8632 | [support@antibodies-online.com](mailto:support@antibodies-online.com)

## Product Details

---

Purification:	The DNAs were purified using PowerPrep HP Plasmid isolation kits for transfection ready plasmids.
Sequencing Primer:	VP1.5 (forward) 5'GGACTTTCCTCAAATGTTCG 3', XL39 (reverse) 5'ATTAGGACAAGGCTGGTGGG 3'
Components:	<ul style="list-style-type: none"><li>• The cDNA clone is shipped in a 2-D bar-coded Matrix tube as dried plasmid DNA.</li><li>• The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.</li></ul>

## Target Details

---

Gene:	KIR3DP1
Alternative Name:	KIR3DP1
Background:	<p>Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several 'framework' genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules, thus, KIR proteins are thought to play an important role in regulation of the immune response. This gene is one of the 'framework' loci that is present on all haplotypes. This locus represents an alternate copy of KIR3DP1 that is represented in a small percentage of the population and may encode a functional protein. The other copy is considered to be a pseudogene. [provided by RefSeq].</p>
NCBI Accession:	<a href="#">NM_001015070</a> , <a href="#">NP_001015070</a>

## Application Details

---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

---

Format: Lyophilized

Storage: RT,-20 °C

Storage Comment: The lyophilized plasmid is stable for up to one year when stored at ambient temperature. Following dissolution in 100 µL dH<sub>2</sub>O, store at -20 °C. Lyophilized primers are stable for up to one year when stored at ambient temperature. Following dissolution in 10 µL dH<sub>2</sub>O, store at -20 °C.

Expiry Date: 12 months

## Publications

---

Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (1991)