## -online.com genomics

## Datasheet for ABIN3376624 Human KIR2DL5B cDNA Clone in Mammalian Expression Vector

Overview	
Quantity:	10 µg
Gene:	KIR2DL5B
Species:	Human
Insert:	cDNA
Vector:	Mammalian Expression Vector
Application:	Protein Expression (PExp)
Product Details	
Purpose:	Untagged full-length cDNA clone from Human KIR2DL5B is ideal for over-expression of native protein for functional studies.
Brand:	TrueClones®
Insert Length:	1100 bp
Vector Backbone:	pCMV6-Neo
Promoter:	Enhanced CMV Promoter, T7 Promoter
Selectable Marker:	Neomycin
Bacterial Resistance:	Ampicillin
Expression Type:	Transient, Stable
Characteristics:	<ul> <li>These cDNA clones are isolated from full-length cDNA libraries and usually contain the coding sequence as well as the untranslated regions (UTRs) of the mRNA transcript appropriate to the library from which they were isolated.</li> <li>These cDNA clones are ideal for over-expression of native proteins for functional studies. Provided as 10 µg transfection-ready plasmids.</li> <li>Every lot of primer is tested to provide clean sequencing of cDNA clones.</li> </ul>
Purification:	The DNAs were purified using PowerPrep HP Plasmid isolation kits for transfection ready

## Product Details

	plasmids.
Sequencing Primer:	VP1.5 (forward) 5'GGACTTTCCAAAATGTCG 3', XL39 (reverse) 5'ATTAGGACAAGGCTGGTGGG 3'
Components:	<ul> <li>The cDNA clone is shipped in a 2-D bar-coded Matrix tube as dried plasmid DNA.</li> <li>The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.</li> </ul>

## Target Details

Gene:	KIR2DL5B
Alternative Name:	KIR2DL5B (KIR2DL5B Products)
Background:	Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several 'framework' genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules, thus, KIR proteins are thought to play an important role in regulation of the immune response. [provided by RefSeq, Jul 2008].
NCBI Accession:	NM_001018081, NP_001018091
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Storage:	RT,-20 °C
Storage Comment:	The lyophilized plasmid is stable for up to one year when stored at ambient temperature.

	Following dissolution in 100 $\mu$ L dH2O, store at -20 °C. Lyophilized primers are stable for up to one year when stored at ambient temperature. Following dissolution in 10 $\mu$ L dH2O, store at -20 °C.
Expiry Date:	12 months
Publications	
Product cited in:	Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, ( 1991)