## -online.com genomics





## **Human PKM cDNA Clone in Mammalian Expression Vector**

Overview	
Quantity:	10 μg
Gene:	PKM
Species:	Human
Insert:	cDNA
Vector:	Mammalian Expression Vector
Application:	Protein Expression (PExp)
Product Details	
Purpose:	Untagged full-length cDNA clone from Human PKM is ideal for over-expression of native protein for functional studies.
Brand:	TrueClones®
Insert Length:	2300 bp
Vector Backbone:	pCMV6-XL5
Promoter:	Enhanced CMV Promoter, T7 Promoter
Bacterial Resistance:	Ampicillin
Expression Type:	Transient
Characteristics:	<ul> <li>These cDNA clones are isolated from full-length cDNA libraries and usually contain the coding sequence as well as the untranslated regions (UTRs) of the mRNA transcript appropriate to the library from which they were isolated.</li> <li>These cDNA clones are ideal for over-expression of native proteins for functional studies. Provided as 10 µg transfection-ready plasmids.</li> <li>Every lot of primer is tested to provide clean sequencing of cDNA clones.</li> </ul>
Purification:	The DNAs were purified using PowerPrep HP Plasmid isolation kits for transfection ready plasmids.

## **Product Details** Sequencing Primer: VP1.5 (forward) 5'GGACTTTCCAAAATGTCG 3', XL39 (reverse) 5'ATTAGGACAAGGCTGGTGGG 3' Components: · The cDNA clone is shipped in a 2-D bar-coded Matrix tube as dried plasmid DNA. • The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials. **Target Details PKM** Gene: Alternative Name: PKM (PKM Products) Background: This gene encodes a protein involved in glycolysis. The encoded protein is a pyruvate kinase that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP, generating ATP and pyruvate. This protein has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. This protein has been found to bind Opa protein, a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells, suggesting a role of this protein in bacterial pathogenesis. Several alternatively spliced transcript variants encoding a few distinct isoforms have been reported. [provided by RefSeq, May 2011]. Transcript Variant: This variant (2) differs in the 5' UTR and coding sequence compared to variant 4. The resulting isoform (b, also called M1) is shorter at the N-terminus compared to isoform c. NCBI Accession: NM\_182470, NP\_872270 **Application Details** For Research Use only Restrictions: Handling Format: Lyophilized RT,-20 °C Storage: Storage Comment: The lyophilized plasmid is stable for up to one year when stored at ambient temperature. Following dissolution in 100 µL dH20, store at -20 °C. Lyophilized primers are stable for up to one year when stored at ambient temperature. Following dissolution in 10 µL dH20, store at -20 °C.

12 months

**Expiry Date:** 

## **Publications**

Product	cited	in:

Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (1991)