-online.com genomics

Datasheet for ABIN4632995 Human OR6K2 cDNA Clone in Bacterial Expression Vector (His tag)

Overview

Quantity:	500 ng
Gene:	OR6K2
Species:	Human
Fusion tag:	His tag
Insert:	cDNA
Vector:	Bacterial Expression Vector
Application:	Cloning (Clon)

Product Details

Purpose:	Bacterial expression of Human OR6K2 with His tag
Insert Length:	975 bp
Vector Backbone:	pPB-N-His
Promoter:	T7 Promoter
Bacterial Resistance:	Kanamycin
Expression Type:	Transient
Specificity:	5-Nhel and 3-Xhol Fusion tag: A singel N-terminal 6X-Histidine tag which is cleavable with Thrombin (Size 2.3 kDa)
Sequencing Primer:	T7 promoter primer: 5'-TAATACGACTCACTATAGGG-3', T7 terminator primer: 5'- GCTAGTTATTGCTCAGCGG-3'

Target Details

Gene:	OR6K2
Alternative Name:	OR6K2 (OR6K2 Products)

Order at www.genomics-online.com

USA & Canada: +1 877 302 8632 | support@antibodies-online.com

Page 1/2 | Product datasheet for ABIN4632995 | 10/07/2023 | Copyright antibodies-online. All rights reserved.

Target	Details
--------	---------

NCBI Accession:

NM_001005279

Application Details

Application Notes:	The pPB vectors are low-medium copy number vectors in which the gene expression is driven
	by the strong T7 promoter.
	Below are some basic guidelines for using the pPB vectors for protein production:
	1. The pPB vectors are designed to be used with E. coli strains that are DE3 lysogens i.e. the
	host E. coli cell has a source of T7 RNA polymerase.
	2. Recombinant protein induction is usually done at OD600 of 0.6-1.2 using Isopropyl β -D-1-
	thiogalactopyranoside (IPTG) at a final concentration of 0.05 -1mM.
	3. The ideal concentration of IPTG must be determined empirically for each recombinant
	protein/cell-line. Similarly, the length of time and temperature for induction provide other
	variables that need to be optimized on a case-to-case basis.
	4. For toxic proteins, it is recommended to go for shorter induction time and also to try and
	suppress basal recombinant gene expression through (a) addition of glucose or use of pLysS
	plasmid. Please note that special cell-lines are also available in the market that cater to
	expression of toxic proteins.
	5. Once grown for the desired length of time, harvest cells by centrifugation and either freeze
	the cells at -80°C (as such or after re-suspending in the desired buffer) or proceed with the
	purification.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	10 mM Tris-HCI, 1 mM EDTA, pH 8.0
Storage:	-20 °C
Storage Comment:	1 year when stored at -20° C or lower in a non-frost free freezer.
Expiry Date:	12 months
Publications	
Product cited in:	Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (
	1991)