

Datasheet for ABIN4835015

Human MAPKAPK2 cDNA Clone in Bacterial Expression Vector (His-GST)

Overview

| | |
|--------------|-----------------------------|
| Quantity: | 500 ng |
| Gene: | MAPKAP Kinase 2 (MAPKAPK2) |
| Species: | Human |
| Fusion tag: | His-GST |
| Insert: | cDNA |
| Vector: | Bacterial Expression Vector |
| Application: | Cloning (Clon) |

Product Details

| | |
|-----------------------|--|
| Purpose: | Bacterial expression of Human MAPKAPK2 with His-GST |
| Insert Length: | 1203 bp |
| Vector Backbone: | pPB-His-GST |
| Promoter: | T7 Promoter |
| Bacterial Resistance: | Kanamycin |
| Expression Type: | Transient |
| Specificity: | 5-NheI and 3-XhoI Fusion tag: Dual N-terminal tag, 6X Histidine followed by Glutathione-S-Transferase Protein which is cleavable with TEV (Size 27.9 kDa) |
| Sequencing Primer: | GST Forward primer: 5'-CACGTTTGGTGGTGGCGAC3', T7 terminator primer: 5'-GCTAGTTATTGCTCAGCGG-3' |

Target Details

| | |
|-------|----------------------------|
| Gene: | MAPKAP Kinase 2 (MAPKAPK2) |
|-------|----------------------------|

Order at www.genomics-online.com

USA & Canada: +1 877 302 8632 | support@antibodies-online.com

Target Details

Alternative Name: MAPKAPK2 ([MAPKAPK2 Products](#))

NCBI Accession: [NM_004759](#)

Application Details

Application Notes: The pPB vectors are low-medium copy number vectors in which the gene expression is driven by the strong T7 promoter.

Below are some basic guidelines for using the pPB vectors for protein production:

1. The pPB vectors are designed to be used with E. coli strains that are DE3 lysogens i.e. the host E. coli cell has a source of T7 RNA polymerase.
2. Recombinant protein induction is usually done at OD600 of 0.6-1.2 using Isopropyl β -D-1-thiogalactopyranoside (IPTG) at a final concentration of 0.05 -1mM.
3. The ideal concentration of IPTG must be determined empirically for each recombinant protein/cell-line. Similarly, the length of time and temperature for induction provide other variables that need to be optimized on a case-to-case basis.
4. For toxic proteins, it is recommended to go for shorter induction time and also to try and suppress basal recombinant gene expression through (a) addition of glucose or use of pLysS plasmid. Please note that special cell-lines are also available in the market that cater to expression of toxic proteins.
5. Once grown for the desired length of time, harvest cells by centrifugation and either freeze the cells at -80°C (as such or after re-suspending in the desired buffer) or proceed with the purification.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 10 mM Tris-HCl, 1 mM EDTA, pH 8.0

Storage: -20 °C

Storage Comment: 1 year when stored at -20° C or lower in a non-frost free freezer.

Expiry Date: 12 months

Publications

Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (

Order at www.genomics-online.com

USA & Canada: +1 877 302 8632 | support@antibodies-online.com

1991)