-online.com





Datasheet for ABIN4919262

Human KHDC1L ORF Clone in Mammalian Expression Vector (DYKDDDDK Tag)

Overview	
Quantity:	10 μg
Gene:	KHDC1L
Species:	Human
Fusion tag:	DYKDDDDK Tag
Insert:	ORF
Vector:	Mammalian Expression Vector
Application:	Protein Expression (PExp)
Product Details	
Purpose:	Expression/transfection ready cDNA ORF clone of Human KHDC1L with C terminal DYKDDDDK
	tag is ideal for express proteins in E.coli & mammalian cells.
Brand:	GenEZ™
Insert Length:	387 bp
Vector Backbone:	pcDNA3.1+C-(K)-DYK
Promoter:	CMV Promoter
Selectable Marker:	Neomycin
Bacterial Resistance:	Ampicillin
Expression Type:	Transient, Stable
Sequence:	ATGGCCGTGG GAACGAGTGC TCTCAGCAAG GAGCCGTGGT GGACCCTGCC CGAAAACTTT
	CATTCTCCAA TGGTGTTCCA CATGGAAGAG GACCAGGAGG AGCTCATCTT CGGACTTGAT
	GACACGTACC TTCGCTGCAT TGAGCTGCAC AGCCACACCC TTATTCAGCT GGAGAGGTGT
	TTCACAGCTA CAGGCCAGAC ACGTGTGACT GTAGTCGGAC CACCAATGGC AAAGCAGTGG
	CTGCTGCTCA TGTTCCATTG CGTGGGGAGC CAGGACTCCA AGTGTCACGC TCGAGGTCTG

Product Details

- Toddot Betaile	
	AAGATGCTAG AGCGTGTCCG AAGCCAGCCC CTGACCAATG ATGACCTGGT CACCTCCGTT
	AGCCTGCCAC CGTACACCGG AGACTGA
Specificity:	ORF Insert Method: CloneEZ® Seamless cloning technology, recombination-based cloning
	technology
Characteristics:	Gene cDNA ORF clone sequences were retrieved from the NCBI Reference Sequence Database
	(RefSeq). These sequences represent the protein coding region of the gene cDNA ORF which is
	encoded by the open reading frame (ORF) sequence.
Sequencing Primer:	Forward primer: 5'-TAATACGACTCACTATAGGG-3'
	Reverse primer: 5'-CCTCGACTGTGCCTTCTA-3'
Grade:	End-sequenced
Components:	The GenEZ ORF clone is delivered as 10 µg of lyophilized plasmid DNA in a vial.
Target Details	
Gene:	KHDC1L
Alternative Name:	KHDC1L (KHDC1L Products)
Gene ID:	100129128
NCBI Accession:	NM_001126063
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Storage:	RT/-20 °C
Storage Comment:	Keep the vial sealed and store at -20°C for long-term storage.
	 Before use, centrifuge the vial at 6,000 g x g for 1 minute at 4°C. Open the lid and add 100 µl (or other volume depending on your desired final concentration)
	of distilled water (or TE buffer) to dissolve the DNA.
	• If necessary, heat the solution at 50°C for 15 minutes to dissolve the DNA.
	 Close the lid and vortex the vial for 1 minute. Aliquot the dissolved plasmid DNA and store in small aliquots at -20°C.
	August the dissolved plastific bigh and stole in strial aliquots at 20 c.

Handling	
Expiry Date:	12 months
Publications	
Product cited in:	Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (
	1991)