# -online.com genomics

## Datasheet for ABIN4920872

Human ZNF717 ORF Clone in Mammalian Expression Vector (DYKDDDDK Tag)

Overview	
Quantity:	10 µg
Gene:	ZNF717
Species:	Human
Fusion tag:	DYKDDDDK Tag
Insert:	ORF
Vector:	Mammalian Expression Vector
Application:	Protein Expression (PExp)
Product Details	
Purpose:	Expression/transfection ready cDNA ORF clone of Human ZNF717 with C terminal DYKDDDDK
	tag is ideal for express proteins in E.coli & mammalian cells.
Brand:	GenEZ™
Insert Length:	345 bp
Vector Backbone:	pcDNA3.1+C-(K)-DYK
Promoter:	CMV Promoter
Selectable Marker:	Neomycin
Bacterial Resistance:	Ampicillin
Expression Type:	Transient, Stable
Sequence:	ATGTTTCCAG TGTTCTCTGG CTGTTTCCAA GAGCTACAAG AAAAGAATAA ATCTCTGGAG
	TTGGTGTCCT TTGAGGAGGT AGCTGTGCAC TTCACCTGGG AGGAGTGGCA GGACCTGGAT
	GATGCTCAGA GGACCCTGTA CAGGGACGTG ATGCTGGAGA CCTACAGCAG CCTGGTATCA
	TTGGGGCATT ACATTACCAA ACCTGAGATG ATCTTCAAGC TAGAGCAAGG AGCAGAGCCA
	TGGATAGTAG AAGAAACCCC AAACCTGAGA CTTTCAGGTG GAAGCAAGAA GCAAGTTTTC

#### Product Details

	TCAGGTATTT GCCACAGGAG CCTGGTGGAG CTCCAGGAGG TTTGA
Specificity:	ORF Insert Method: CloneEZ® Seamless cloning technology, recombination-based cloning technology
Characteristics:	Gene cDNA ORF clone sequences were retrieved from the NCBI Reference Sequence Database (RefSeq). These sequences represent the protein coding region of the gene cDNA ORF which is encoded by the open reading frame (ORF) sequence.
Sequencing Primer:	<ul> <li>Forward primer: 5'-TAATACGACTCACTATAGGG-3'</li> <li>Reverse primer: 5'-CCTCGACTGTGCCTTCTA-3'</li> </ul>
Grade:	End-sequenced
Components:	The GenEZ ORF clone is delivered as 10 $\mu g$ of lyophilized plasmid DNA in a vial.
Target Details	
Gene:	ZNF717
Alternative Name:	ZNF717 (ZNF717 Products)
Gene ID:	100131827
NCBI Accession:	NM_001290210

### Application Details

Restrictions: For Research Use only

#### Handling

Format:	Lyophilized
Storage:	RT/-20 °C
Storage Comment:	<ul> <li>Keep the vial sealed and store at -20°C for long-term storage.</li> </ul>
	• Before use, centrifuge the vial at 6,000 g x g for 1 minute at $4^{\circ}$ C.
	• Open the lid and add 100 $\mu$ l (or other volume depending on your desired final concentration)
	of distilled water (or TE buffer) to dissolve the DNA.
	<ul> <li>If necessary, heat the solution at 50°C for 15 minutes to dissolve the DNA.</li> </ul>
	Close the lid and vortex the vial for 1 minute.
	<ul> <li>Aliquot the dissolved plasmid DNA and store in small aliquots at -20°C.</li> </ul>

Expiry Date:

Publications	
Product cited in:	Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (
	1991)