

Datasheet for ABIN4925275

Human POLR2J3 ORF Clone in Mammalian Expression Vector (DYKDDDDK Tag)

Overview

| | |
|--------------|-----------------------------|
| Quantity: | 10 µg |
| Gene: | POLR2J3 |
| Species: | Human |
| Fusion tag: | DYKDDDDK Tag |
| Insert: | ORF |
| Vector: | Mammalian Expression Vector |
| Application: | Protein Expression (PEXP) |

Product Details

| | |
|-----------------------|--|
| Purpose: | Expression/transfection ready cDNA ORF clone of Human POLR2J3 with C terminal DYKDDDDK tag is ideal for express proteins in E.coli & mammalian cells. |
| Brand: | GenEZ™ |
| Insert Length: | 348 bp |
| Vector Backbone: | pcDNA3.1+C-(K)-DYK |
| Promoter: | CMV Promoter |
| Selectable Marker: | Neomycin |
| Bacterial Resistance: | Ampicillin |
| Expression Type: | Transient, Stable |
| Sequence: | ATGAACGCCC CTCCAGCCTT CGAGTCGTTT TTGCTCTTCG AGGGCGAGAA GATCACCATT AACAAGGACA CCAAGGTACC CAATGCCTGT TTATTCACCA TCAACAAAGA AGACCACACA CTGGGAAACA TCATTAAATC ACAACTCCTA AAAGACCCGC AAGTGCTATT TGCTGGCTAC AAAGTCCCCC ACCCCTTGGA GCACAAGATC ATCATCCGAG TGCAGACCAC GCCGGACTAC |

Order at www.genomics-online.com

USA & Canada: +1 877 302 8632 | support@antibodies-online.com

Product Details

AGCCCCCAGG AAGCCTTTAC CAACGCCATC ACCGACCTCA TCAGTGAGCT GTCCCTGCTG
GAGGAGCGCT TCCGGACGTG CCTGCTTCCC CTTCGCCTTC TGCCGTGA

Specificity: ORF Insert Method: CloneEZ® Seamless cloning technology, recombination-based cloning technology

Characteristics: Gene cDNA ORF clone sequences were retrieved from the NCBI Reference Sequence Database (RefSeq). These sequences represent the protein coding region of the gene cDNA ORF which is encoded by the open reading frame (ORF) sequence.

Sequencing Primer:

- Forward primer: 5'-TAATACGACTCACTATAGGG-3'
- Reverse primer: 5'-CCTCGACTGTGCCTTCTA-3'

Grade: End-sequenced

Components: The GenEZ ORF clone is delivered as 10 µg of lyophilized plasmid DNA in a vial.

Target Details

Gene: POLR2J3

Alternative Name: POLR2J3 ([POLR2J3 Products](#))

Background: This gene is a member of the RNA polymerase II subunit 11 gene family, which includes three genes in a cluster on chromosome 7q22.1 and a pseudogene on chromosome 7p13. The founding member of this family, DNA directed RNA polymerase II polypeptide J, has been shown to encode a subunit of RNA polymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. This locus produces multiple, alternatively spliced transcripts that potentially express isoforms with distinct C-termini compared to DNA directed RNA polymerase II polypeptide J. Most or all variants are spliced to include additional non-coding exons at the 3' end which makes them candidates for nonsense-mediated decay (NMD). Consequently, it is not known if this locus expresses a protein or proteins in vivo. [provided by RefSeq, Jul 2008].

Gene ID: 548644

NCBI Accession: [NM_001097615](#)

Application Details

Restrictions: For Research Use only

Order at www.genomics-online.com

USA & Canada: +1 877 302 8632 | support@antibodies-online.com

Handling

Format: Lyophilized

Storage: RT/-20 °C

Storage Comment:

- Keep the vial sealed and store at -20°C for long-term storage.
- Before use, centrifuge the vial at 6,000 g x g for 1 minute at 4°C.
- Open the lid and add 100 µl (or other volume depending on your desired final concentration) of distilled water (or TE buffer) to dissolve the DNA.
- If necessary, heat the solution at 50°C for 15 minutes to dissolve the DNA.
- Close the lid and vortex the vial for 1 minute.
- Aliquot the dissolved plasmid DNA and store in small aliquots at -20°C.

Expiry Date: 12 months

Publications

Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (1991)