

Datasheet for ABIN5328231

Human APOBEC3A_B ORF Clone in Mammalian Expression Vector

Overview

Quantity:	10 µg
Gene:	APOBEC3A/B (APOBEC3A_B)
Species:	Human
Fusion tag:	Myc-DYKDDDDK Tag
Insert:	ORF
Vector:	Mammalian Expression Vector
Application:	Protein Expression (PEXP)

Product Details

Purpose:	Mammalian Vector with ORF clone of Human APOBEC3A and APOBEC3B deletion hybrid (APOBEC3A_B)
Brand:	TrueORF
Insert Length:	600 bp
Vector Backbone:	pCMV6-Entry
Promoter:	CMV Promoter
Bacterial Resistance:	Kanamycin
Expression Type:	Transient
Specificity:	Restriction Site: SgfI-MluI
Sequencing Primer:	VP1.5 (forward) 5'GGACTTTC AAAATGTTCG 3', XL39 (reverse) 5'ATTAGGACAAGGCTGGTGGG 3'
Grade:	End-sequenced
Components:	The ORF clone is ion-exchange column purified, transfection-ready dried plasmid DNA, and shipped with 2 vector sequencing primers.

Order at www.genomics-online.com

USA & Canada: +1 877 302 8632 | support@antibodies-online.com

Target Details

Gene: [APOBEC3A/B \(APOBEC3A_B\)](#)

Abstract: [APOBEC3A_B Products](#)

Background: This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster, thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. The protein encoded by this gene lacks the zinc binding activity of other family members. The protein plays a role in immunity, by restricting transmission of foreign DNA such as viruses. One mechanism of foreign DNA restriction is deamination of foreign double-stranded DNA cytidines to uridines, which leads to DNA degradation. However, other mechanisms are also thought to be involved, as anti-viral effect is not dependent on deaminase activity. The protein encoded by this gene is the same as that encoded by APOBEC3A, however, this gene is a hybrid gene that results from the deletion of approximately 29.5 kb of sequence between the APOBEC3A gene and the adjacent gene APOBEC3B. The breakpoints of the deletion are within the two genes, so the deletion hybrid is predicted to have the promoter and coding region of APOBEC3A, but the 3' UTR of APOBEC3B.

NCBI Accession: [NM_001193289](#), [NP_001180218](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Storage: 4 °C/-20 °C

Publications

Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (1991)