## -online.com genomics



Datasheet for ABIN5480600

## Human OR9I1 ORF Clone in Mammalian Expression Vector (Myc-DYKDDDDK Tag)

Overview	
Quantity:	10 μg
Gene:	OR9I1
Species:	Human
Fusion tag:	Myc-DYKDDDDK Tag
Insert:	ORF
Vector:	Mammalian Expression Vector
Application:	Protein Expression (PExp)
Product Details	
Purpose:	Mammalian Vector with ORF clone of Human olfactory receptor, family 9, subfamily I, member
	1 (OR9I1)
Brand:	TrueORF
Insert Length:	945 bp
Vector Backbone:	pCMV6-Entry
Promoter:	CMV Promoter
Bacterial Resistance:	Kanamycin
Expression Type:	Transient
Specificity:	Restriction Site: Sgfl-Mlul
Sequencing Primer:	VP1.5 (forward) 5'GGACTTTCCAAAATGTCG 3', XL39 (reverse) 5'ATTAGGACAAGGCTGGTGGG 3'
Grade:	End-sequenced
Components:	The ORF clone is ion-exchange column purified, transfection-ready dried plasmid DNA, and

shipped with 2 vector sequencing primers.

## Target Details

that triggers the perception of a smell. The olfactory receptor proteins are members of a family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Ol receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for the organism is independent of other organisms.  NCBI Accession:  NML001005211, NP_001005211  Application Details  Restrictions:  For Research Use only  Handling  Format:  Lyophilized  Storage:  4 °C/-20 °C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3		
Background:  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal resthat triggers the perception of a smell. The olfactory receptor proteins are members of a family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Ol receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for the organism is independent of other organisms.  NCBI Accession:  NM_001005211, NP_001005211  Application Details  Restrictions:  For Research Use only  Handling  Format:  Lyophilized  Storage:  4 *C/-20 *C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3	Gene:	OR9I1
that triggers the perception of a smell. The olfactory receptor proteins are members of a family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Ol receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for the organism is independent of other organisms.  NCBI Accession:  NML001005211, NP_001005211  Application Details  Restrictions:  For Research Use only  Handling  Format:  Lyophilized  Storage:  4 °C/-20 °C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3	Abstract:	OR9I1 Products
family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Of receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for the organism is independent of other organisms.  NCBI Accession:  NM_001005211, NP_001005211  Application Details  Restrictions:  For Research Use only  Handling  Format:  Lyophilized  Storage:  4 °C/-20 °C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-5	Background:	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response
receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for the organism is independent of other organisms.  NCBI Accession:  NM_001005211, NP_001005211  Application Details  Restrictions:  For Research Use only  Handling  Format:  Lyophilized  Storage:  4 °C/-20 °C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3		that triggers the perception of a smell. The olfactory receptor proteins are members of a large
hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for the organism is independent of other organisms.  NCBI Accession:  NM_001005211, NP_001005211  Application Details  Restrictions:  For Research Use only  Handling  Format:  Lyophilized  Storage:  4 °C/-20 °C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3		family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory
transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for the organism is independent of other organisms.  NCBI Accession:  NM_001005211, NP_001005211  Application Details  Restrictions:  For Research Use only  Handling  Format:  Lyophilized  Storage:  4 *C/-20 *C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3		receptors share a 7-transmembrane domain structure with many neurotransmitter and
genome. The nomenclature assigned to the olfactory receptor genes and proteins for the organism is independent of other organisms.  NCBI Accession:  NM_001005211, NP_001005211  Application Details  Restrictions:  For Research Use only  Handling  Format:  Lyophilized  Storage:  4 °C/-20 °C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3		hormone receptors and are responsible for the recognition and G protein-mediated
organism is independent of other organisms.  NCBI Accession:  NM_001005211, NP_001005211  Application Details  Restrictions:  For Research Use only  Handling  Format:  Lyophilized  Storage:  4 °C/-20 °C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3		transduction of odorant signals. The olfactory receptor gene family is the largest in the
NCBI Accession:  NM_001005211, NP_001005211  Application Details  Restrictions:  For Research Use only  Handling  Format:  Lyophilized  Storage:  4 °C/-20 °C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in:, Vol. 1363, Issue Nucleic acids research, pp. 28-3		genome. The nomenclature assigned to the olfactory receptor genes and proteins for this
Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Storage: 4 °C/-20 °C  Publications  Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3		organism is independent of other organisms.
Restrictions: For Research Use only  Handling  Format: Lyophilized  Storage: 4 °C/-20 °C  Publications  Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3	NCBI Accession:	NM_001005211, NP_001005211
Format: Lyophilized  Storage: 4 °C/-20 °C  Publications  Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3	Application Details	
Format:  Lyophilized  Storage:  4 °C/-20 °C  Publications  Product cited in:  Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3	Restrictions:	For Research Use only
Storage: 4 °C/-20 °C  Publications  Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3	Handling	
Publications  Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3	Format:	Lyophilized
Product cited in: Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-3	Storage:	4 °C/-20 °C
	Publications	
1001)	Product cited in:	Johnson, Drugan, Miller, Evans: "38" in: , Vol. 1363, Issue Nucleic acids research, pp. 28-39, (
1991)		1991)